



Children's Urology Group, PL

DENNIS L. HOOVER, M.D. • E. MICHAEL REISMAN, M.D.
MARK E. KOLLIGIAN, M.D. • YVES L. HOMSY, M.D.
ETHAN G. POLSKY, M.D.

Certified American Board of Urology

(813) 874-7500, (727) 456-1055

www.childreurologygroup.com

MOGEN CLAMP

NEWBORN CIRCUMCISION CARE AFTER NEWBORN CIRCUMCISION

Your son's circumcision was performed using the "Mogen Clamp" technique. The following instructions should be followed to ensure proper care following the procedure.

BANDAGE/DRESSING: Remove the bandage after your son first urinates. Apply large amounts of antibiotic ointment (Neosporin/Bacitracin/Triple antibiotic).

BATHING: On the first day following the circumcision, bathe your son in a warm water bath (**no soap**) 3 - 4 times per day. Apply very generous amounts of the antibiotic ointment after the baths and allow the penis to "air dry".

CLEANING: You may clean him with diaper wipes or soap and water at diaper changes. You may apply an over the counter antibiotic ointment to the tip of the penis (Bacitracin, Neosporin, Triple antibiotic), especially if the penis tends to stick to the diaper. Please **DO NOT** apply Vaseline to the penis. Typically a bandage is not necessary.

IRRITABILITY: Your child may be irritable during the first 24 hours following the circumcision. Cuddling and feeding may console him, just as cuddling and feeding calmed him immediately after the circumcision was performed. Tylenol may be given (as needed every 4 hours). **Avoid** Ibuprofen (Motrin, Advil).

BLEEDING and SWELLING: Blood spotting on the diaper is expected. Moderate swelling and redness of the head of the penis is expected. If you notice a continuous ooze of blood, apply pressure for 10 minutes. If bleeding persists, please call the office. If you notice redness around the shaft of the penis or fever greater than 101.5F, please call the office. It is not unusual to have a bruise at the base of the penis if local anesthesia was injected. This will go away.

POSTOPERATIVE OFFICE VISIT: Your doctor may want you to schedule a postoperative visit for 2 weeks following the date of surgery by calling the nearest office during the hours of 8:00 a.m. to 5:00 p.m., Monday through Friday.

St. Petersburg: 727.456.1055

Tampa: 813.874.7500

To reach the doctor after the above hours and on weekends, the answering service will direct you from the above telephone numbers.

Infants' Acetaminophen Concentrated Drops – "Old" Concentration			Infants' Acetaminophen Oral Suspension – "New" Concentration		
80 mg/0.8 mL			160 mg/5mL		
Weight (lb)	Age (mos)	Dose (mL)	Weight (lb)	Age (mos)	Dose (mL)
6-11 lbs	0-3 mos	0.4 mL	6-11 lbs	0-3 mos	1.25 mL
12-17 lbs	4-11 mos	0.8 mL	12-17 lbs	4-11 mos	2.5 mL
18-23 lbs	12-23 mos	1.2 mL	18-23 lbs	12-23 mos	3.75 mL
24-35 lbs	24-36 mos	1.6 mL	24-35 lbs	24-36 mos	5 mL